SECTION K

1. NARRATIVE OF

-

4. INTRA-PROCEDURE ANIMAL

TABLE 5.A. INTRA-PROCEDURE ANESTHESIA OR CHEMICAL RESTRAINT

To add additional anesthetic agents, please attach a separate document

ANESTHETIC AGENT	DOSE	ROUTE OF ADMINISTRATION	FREQUENCY OF ADMINISRATION	DURATION OF TREATMENT

6. NEUROMUSCULAR BLOCKING AGENTS (PARALYTICS)

Will neuromuscular blocking agents (paralytics) be used at any time during the procedure?

☐ NO. Neuromuscular blocking agents will not be used for the procedure.
☐ YES. Neuromuscular blocking agents will be used. (Provide details below)

TABLE 6.A. PARALYTICS

To add additional paralytic agents, please attach a separate document.

7. MONITORING DEPTH OF ANESTHESIA DURING PROCEDURES

- ☐ Body temperature
- ☐ Oxygen saturation
- ☐ Capillary Refill Time
- ☐ EKG
- ☐ Reflex (specify):
- ☐ Other (specify):
- ☐ Other (specify):

Specify the frequency at which the above indices will be recorded:

10. PAIN MANAGEMENT INTRA OR POST-PROCEDURE

NOTE: The IACUC encourages the use of pre-emptive analgesia for pain management. Analgesia should be provided as early as possible in the procedure if it is expected to be painful or result in residual pain, ideally before the procedure begins.

Is the procedure expected to cause pain or result in residual pain?

☐ Yes ☐ No

If pain is expected, will analgesia be provided for pain relief?

☐ NO. Analgesia will no

12. Humane Endpoints That Will Prompt Intervention to Prevent Continued Pain or Distress

According to The Guide, information that is critical to the IACUC's assessment of appropriate endpoint consideration within a protocol includes precisely defining the humane endpoint (including assessment criteria); the frequency of animal observation; training of personnel responsible for assessment and recognition of the humane endpoint; and the response required upon reaching the humane endpoint. The IACUC has determined that the list below defines the commonly accepted clinical milestones which should be regarded as humane endpoints for most terrestrial animal studies. Choose all of those which are appropriate for the species being used. For each milestone, indicate the action that will be taken. Add other milestones (in the row marked 'other') if applicable for defining the humane endpoints for the proposed study.

CLINICAL OBSERVATION/

r4 Tm [-1.22 Td [(t)-4 (BT 0 scn 125 Tc -0l)10 (th)10 (e)3 >>BDC 1 scn 74.52 30214.64 re

CLINICAL OBSERVATION MILESTONE	APPLICABLE TO MY PROPOSAL	FREQUENCY OF OBSERVATIO (e.g., 4 hrs., 12 hrs., weekly)	PROTOCOL PERSONNEL ARE TRAINED TO RECOGNIZE	RESPONSE REQUIRED UPON REACHING THE HUMANE ENDPOINT	PROVIDE DURATION (DAYS, WEEKS, ETC.) OF MONITORING OR A SCIENTIFIC JUSTIFICATION FOR NOT USING THE MILESTONES LISTED
(will require individual housing to effectively assess)				☐	

CLINICAL OBSERVATION MILESTONE	APPLICABLE TO MY PROPOSAL	FREQUENCY OF OBSERVATIO (e.g., 4 hrs., 12 hrs., weekly)	PROTOCOL PERSONNEL ARE TRAINED TO RECOGNIZE	RESPONSE REQUIRED UPON REACHING THE HUMANE ENDPOINT	PROVIDE DURATION (DAYS, WEEKS, ETC.) OF MONITORING OR A SCIENTIFIC JUSTIFICATION FOR NOT USING THE MILESTONES LISTED
Poor posture or ambulating difficulty (Ex: tense, tucked-up, stiff gait)	☐yes☐ no		☐yes☐ no	☐Consult Vet ☐ Euthanize	
Lost hair coat condition (Ex: ruffled fur, lack of grooming, piloerection)	☐yes☐ no		☐yes☐ no	☐Consult Vet ☐ Euthanize	
Sudden activity level change (Ex: restlessness, pacing, reluctance to move)	☐yes☐ no		☐yes☐ no	☐Consult Vet ☐ Euthanize	
Unexpected sweating or salivation (Ex: stressed rodents salivate excessively when stressed)	☐yes☐ no		☐yes☐ no	☐Consult Vet ☐ Euthanize	

CLINICAL OBSERVATION MILESTONE	APPLICABLE TO MY PROPOSAL	FREQUENCY OF OBSERVATIO (e.g., 4 hrs., 12 hrs., weekly)	PROTOCOL PERSONNEL ARE TRAINED TO RECOGNIZE	RESPONSE REQUIRED

CLINICAL OBSERVATION MILESTONE	APPLICABLE TO MY PROPOSAL	FREQUENCY OF OBSERVATIO (e.g., 4 hrs., 12 hrs., weekly)	PROTOCOL PERSONNEL ARE TRAINED TO RECOGNIZE	RESPONSE REQUIRED UPON REACHING THE HUMANE ENDPOINT	PROVIDE DURATION (DAYS, WEEKS, ETC.) OF