The Health and Safety Effects of Accidents on Intermodal Transportation Workers: A Study of Psychological Health Concerns and Depression of Operating Employees

DISCLAIMER

The contents of this report reflect the views of the authors, who are responsible for the facts and the accuracy of the information presented herein. This document is disseminated under the sponsorship of the Department of Transportation University Transportation Centers Program, in the interest of information exchange. The U.S. Government assumes no liability for the contents or use thereof.

Table of Contents

Introduction 6
Definition of Depression 6
Symptoms of Depression 6
Prevalence of Depression 8
Costs of Depression 9
Purpose 9
Methodology 10
Participants 10
Measures 15
CESD 15
Beck Depression Inventory (BDI-II) 16
Single Item Measures 17
Results 17
Prevalence of Depression 17
BDI 17
CESD 19
CESD Short Form 22
Single Item Measures of Depression 24
Relationships with Other Variables 26
Age 26
Suicide 27
Concentration 28
Fatigue 29
Injuries 31
Safety at Work 31
Absenteeism 31
Critical Incidents 32

List of Tables

Table 1. Locations and Sample Sizes. 10
Table 2. Demographic Characteristics of Freight Samples 11
Table 3. Demographic Characteristics of Transit Sample 12
Table 4. Major Depression Likely 19
Table 5. Major Depression Likely – CESD Transit Sample..... 21
Table 6. Frequency Distribution for CESD - 10 Item Short Form 23
Table 7. To what extent have you been feeling unhappy and depressed? 24
Table 8. Single item measure of depression in freight and transit rail samples..... 25
Table 9. Suicidal thoughts or wishes 28
Table 10. BDI - Concentration Difficulty..... 29
Table 11. Fatigue indicators – Sleeping Changes reported by freight rail respondents. 29
Table 12. Fatigue indicators – Tiredness reported by freight rail respondents..... 30

List of Figures

Figure 1. Age Group Distribution for Freight Sample..... 13
F

The Health and Safety Effects of Accidents on Intermodal Transportation Workers: A Study of Psychological Health Concerns and Depression of Operating Employees

Hasin, et al. (2005) reported that the 12-month MDD prevalence rate was 5.28% for the total population and 3.56% for males and 6.87% for females.

working more than 16 years with a mode occurring at 30 to 32 years. This presents an interesting set of demographics when trying to generalize to the entire population.

Table 2. Demographic Characteristics of Freight Samples

Gender		
Male	1062	97.3%

Table 3.



Figure 1. Age Group Distribution for Freight Sample



Figure 2. Age Distribution for Freight Sample



Figure 3. Age Group Distribution for Transit Sample

Measures

CESD

Center for Epidemiological Studies Depression Scale (CES-D) (Radloff, 1977). The CES-D is a 20 item self-report scale designed to assess depressed mood during the past week using a 4 point likert type response scale ranging from 0 "Not at all" to 3 "Most of the time". The CES-D was developed by including items from the BDI, Zung and other previously validated depression measures. It has also been validated with cardiac patients (Penninx et al. 2001) and older populations (Zich et al. 1990) and has good test-retest reliability (Ensel 1986). Scores range from 0 to 60, with higher scores indicating more symptoms of depression. CESD scores of 16 to 26 are considered indicative of mild depression and scores of 27 or more indicative of major depression (Zich et al. 1990, Ensel 1986). Zich, Attkisson & Greenfield (1990) found a cut-off score of 27 more useful for screening medical patients for depression than the usual cut-off score of 16. The cutoffs have been successfully used in studies by Ensel 1986; Zich, Attkisson et al. 1990; Logsdon, McBride et al. 1994; Geisser, Roth et al. 1997.

A study by Pandya, Metz and Patten (2005) with Multiple Sclerosis patients found that a cutoff score of >16 resulted in the correct identification of 74.5% of patients who were DSM-IV diagnosable with a Major Depressive Episode (MDE). The chart below (permission pending) shows that a cutoff score of >=28 will result in the correct classification of more than 85% of persons reporting an MDE. pe[1s/rsons req0(g)1-48(req0(g1-0ilEnse)4(0 4tha)4(s)59(4 TJ)-49(MD

did not meet criteria for MDD (positive predictive value = .28). The authors concluded that the standard cut-score was valid but that a score of 34 yielded a higher specificity (.95) and over

Single Item Measures

Several studies have been conducted that have attempted to shorten the BDI and the CESD even further. More recently there have been several published reports of attempts to use single item measures as screening tools in various setting to facilitate additional screening and early identification of depressive conditions. One item from the General Health Questionnaire (GHQ-12; Goldberg, 1972) specifically asks for information about depression, The GHQ-12 has been used to assess levels of depression, anxiety, sleep disturbance and happiness in the general population. One item was taken from the GHQ-12 and reformatted for use with the railroad population. A Likert response format (1= to a little or no degree to 5 = to a very great degree) was used to be consistent with the other items in the survey. To compare the results to previous studies employing the GHQ a binary or dichotomous response format was derived by recoding the responses into either a no = 0, for responses 1,2,3 on the Likert scale or yes = 1, responses 4 and 5, to permit comparisons with other published data.

Results

Prevalence of Depression

BDI

Depression scores were calculated for the various samples using either the Beck Depression Inventory or the CES-D. We began the study with the CES-D



Figure 5. Frequency Distribution of BDI Scores in Total Sample (N=622).

In Figure 5 the histogram of respondents per each BDI value is presented. The following graph and table were complied using the recommended cutoffs (Steer, Brown, Beck and Sanderson, 2001; Beck et al.;1996) from the validation studies of the BDI. The respondents can be classified into mild, moderate and severely depressed categories. The categories depict a population in which a significant portion of the respondents reported levels of depression at or above the traditional cut-offs recommended.



BDI Category	Frequency	Percent	Cumulative Percent
None	477	77.3	77.3
Mild	67	10.9	88.2
Moderate	54	8.8	96.9
Severe	19	3.1	100.0
Total	617	100.0	

Figure 6. Beck Diagnostic Categories

As can be seen from Figure 6, after removing missing data, 77.3% percent were classified as not reporting any depression, 10.9% indicated mild depression, 8.8% moderate depression, and

Figure 9. CESD in Rail Transit Sample.

CESD Short Form

A more recent study by Irwin et al 1999 looked at the utilization of a short form of the CESD and provided specificity and sensitivity analyses for the various cutoffs. In a sample of 40 depressed patients 39 were correctly identified using a cutoff score of >= 4. Sensitivity, Specificity and Positive Predictive Value were 97%, 84%, and 85% respectively.

Using the =>4 point cutoff on the recalculated 10 item CESD -Short Form it can be seen from the following chart that more than half would meet the criteria for having a depressive disorder.


20
15
10
Frequency
0.00
3.00
6.00
9.00
12.00
CESD - 10 Item Short Form

Figure 10

Single Item Measures of Depression

S



Figure 11. GHQ Single item Scores on feeling depressed.

Table 8. Single item measure of depression in freight and transit rail samples.

How often have you felt depressed? (Transit)

	Frequency	Percent	Valid Percent	Cumulative Percent
Rarely	186	62.4	68.6	68.6
Some	52	17.4	19.2	87.8
Occasionally	24	8.1	8.9	96.7
Most	9	3.0	3.3	100.0
Total	271	90.9	100.0	
Missing	27	9.1		
Total	298	100.0		

How often have you felt depressed? (Freight)

	–	Percent	Valid Percent	Cumulative Percent
Rarely	58	5.0	61.7	61.7
Some	[redacted]	2.0	24.5	86.2
Occasionally	9	.8	9.6	95.7
Most	[redacted]	.3	4.3	100.0
Total				

participants reported feeling depressed. As can be seen in Table 8 a little over 4% of the freight sample felt depressed most of the time as did 3.3% of the Transit sample.

Relationships with Other Variables

There were a number of correlates of the BDI. One of the first things that we needed to determine is whether there was any correlation between the BDI and some of the key demographic variables.

Age

There are some misconceptions about the relationship between age and depression with some thinking that depression increases with age. However, some studies have shown a

In the present study the correlation between the BDI and age was significant (r =-.133, p<..002, N=560) and supported the notion that younger ages generally report more depression. As can be seen in Figure 12, the BDI measure of depression declined gradually with age. In addition, BDI was below clinical cutoffs (cutoff = 14) on the average for all five age groups. Similarly, for the CESD the correlations between age and depression were non-significant in both the Freight and the Transit samples.

Suicide

Suicide has been identified as the eleventh leading cause of death in the U.S. Suicide was the eighth leading cause of death for males and the sixteenth leading cause of death for females in 2004. The overall rate was 10.9 suicide deaths per 100,000 people. (CDC, 2007)_ Major risk factors include: depression and other mental disorders, or a substance-abuse

Results from the suicidal ideation item of the BDI provide interesting data. For example, Item #9 asks respondents to indicate the extent to which they are experiencing suicidal thoughts. As seen in the table below 93.3 percent of respondents were not experiencing any thoughts of killing themselves. However, a relatively small number, 0.5% indicated that they would if they had the chance. While this is a small percent, in a group of 20,000 operating employees 100 people could be actively considering taking their own life. Interestingly, men tend to be more likely to engage in active and violent means of completing suicide than do women who are more likely to use a passive means such as taking pills.

Table 9. Suicidal thoughts or wishes

	Frequency	Percent	Valid Percent	Cumulative Percent
I don't have any thoughts of killing myself	575	49.9	93.3	93.3

they could not concentrate as well as they used to and in fact a combined total of 5.7% report more severe concentration difficulties ("can't keep my mind on anything" and "can't concentrate on anything"). Such difficulties could present some challenges in the operational environment.

Table 10. BDI - Concentration Difficulty

Table 12. Fatigue indicators Tiredness reported by freight rail respondents.

Tiredness or Fatigue

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No more tired than usual	310	26.9	50.2	50.2
	Tired more easily	242	21.0	39.2	89.3
	Too tired to do a lot of the things I used to	51	4.4	8.3	97.6
	Too tired to so most things	15	1.3	2.4	100.0
	Total	618	53.6	100.0	
Missing	System	535	46.4		
Total		1153	100.0		

Two items from the BDI specifically reference fatigue and from the frequency distribution we can see that slightly over 10% of the respondents indicate that they are "too tired to do a lot of the things I used to" and "Too tired to do most things". Furthermore, about 22.4% indicated that they "slept less"; 4.6% reported "sleep less than usual" and 1.1% reported waking and being unable to get back to sleep.

For the N=434 respondents who completed both the BDI and the Epworth in the freight rail sample the correlation between the two scales was statistically significant (r=.315, p<.001). Clearly, there is a moderate and positive relationship between the amount or level of depressed affect or mood and the amount of sleepiness that is reported



Figure 13. Scattergram for BDI and Epworth.

fatigue reported in this particular sample. Moreover, **the relative risk of reporting high levels of fatigue when also reporting depression is 2.49,** meaning that there is almost two and half times the risk of reporting very high levels of fatigue when also experiencing depression.

Results for the CESD in the transit rail sample also show a similar picture. Respondents who completed the CESD and the Epworth produced a significant positive correlation (r = .240, p <.001, N= 275). In effect, the greater the depression the greater the self-reported sleepiness and vice versa.

Injuries

Critical Incidents

Concentration

Results of the analyses indicate that 33% of respondents might not be able to concentrate as well as they used to and in fact a combined total of 5.7% report more severe concentration difficulties ("can't keep my mind on anything" and "can't concentrate on anything"). Such difficulties could present some challenges in the operational environment.

This is an important finding in that, while most people generally associate depression with feelings of sadness and low self-worth they usually don't make the association with concentration. In the rail operating environment concentration, in the face of numerous routine, but high risk or high stakes tasks, can be catastrophic.

The present findings point to the need to inc[(282. Tm[(n.)-26[(n.)] TJET4C>-9-29(inca)] 5( )-9(a)4

can affect the ability of workers to concentrate, attend to their tasks, and experience a satisfying quality of life.

Limitations

There are several limitations of the present study. The most important is the fact that the primary measures used were self-report which provide respondents with the opportunity to self-rate their own subjective experience. These self-ratings have the limitation of being open to bias. In addition the respondents self-selected in responding to the questionnaires.

Another limitation has to do with the fact that the population studies were all gainfully employed and were surveyed when they reported for duty. Despite the adequate response rate (81.3%) it may be the case that this sample might not meet the DSM criteria of "clinically significant distress or impairment in social, occupational or other important areas of function" as required for a DSM-IV diagnosis of Major Depressive Episode.

The sample size upon which the various instruments were administered varies from around six hundred to less. While we had a large sample size additional respondents all completing the same instrument are needed to achieve the highest possible levels of statistical significance.

The use of a structured interview forma to assess the presence of diagnosable symptoms would also have provided greater certainty and confidence in the results. Several structured interviews are available based on the DSM

4. **Public Awareness and Education.** A campaign to alert people to the issues facing employees
5. **Safety and Depression.** Given high levels of depression found in this study employers may also wish to consider the potential impact of depression on safety on operations in the work place. The results how significant correlations between depression safety and concentration. Consequently, employers may wish to more closely monitor levels of depression in their workforce by require. Additional screen, awareness campaigns, publication of National Depression Awareness Week could be a significant first step. Later, more aggressive screening could be done.

References

American Psychiatric Association (APA). (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.

Arnou, R., Meagher, M., Norris, M., & Bramson, R. (2001). Psychometric evaluation of the beck depression inventory-II With Primary Medical Care Medical patients. *Health Psychology*, 20, 112-119.

Ball, R., & Steer, RA. (@003). Mean Beck Depression Inventory –II Scores of outpatients with dysthymic or recurrent episode major depressive disorders. *Psychological Reports*, 93, 507-512.

Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: Guilford Press.

Goldberg, D. P. (1972). *The detection of psychiatric illness by questionnaire* (Maudsley Monograph No. 21). Oxford, England: Oxford University Press.

Hann, D., Winter, K., & Jacobsen, P. (1999). Measurement of depressive symptoms in cancer patients: Evaluation of the CESD Scale. Journal of Psychosomatic Research, 46, 437-443.
Henderson, A.S., Jorm, A.F., Korten, A.E., Jacomb, P., Christensen, H., and Rodgers, B. (1998). Symptoms of depression and anxiety during adult life: evidence for a decline in prevalence with age. Psychological Medicine, 28, 1321-1328.

Ivarsson T, Larsson B, Gillberg C. A 2-4 year follow-up of depressive symptoms, suicidal ideation, and suicide attempts among adolescent psychiatric inpatients. *European Child and Adolescent Psychiatry.* 1998;7:96-104.

K. Fountoulakis, P. Bech, P. Panagiotidis, M. Siamouli, S. Kantartzis, A. Papadopoulou, M. Papadopoulou, S. Kaprinis, E. Kourila, A. Iacovides (200x). Comparison of depressive indices: Reliability, validity, relationship to anxiety and personality and the role of age and life events Journal of Affective Disorders, Volume 97, Issue 1, Pages 187-195

Kessler RC, Borges G, Walters EE: Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. Arch Gen Psychiatry 1999; 56:617–626

Kessler RC; Chiu WT; Demler O; Merikangas KR; Walters EE (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. Arch Gen Psychiatry. 2005 Jun;62 (6):617-27.

Kessler, Hagop, Ames, et (2006) p 156 Am J of Psychiatry

Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, S., Wittchen, H. U., & Kendler, K. S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry , 51*, 8–19.

Kessler, RC, Frank, RG. (1997) the impact of psychiatric disorders on work loss days. Psychological Medicine, 27, 861-873.

Klerman, G. L., Weissman, M. M., Rousaville, B. J., & Sherron, E. S. (1984). *Interpersonal psychotherapy of depression.* New York: Basic Books.

Kogan, E., Kabacoff, R., Hersen, M., & van Hasselt, V. (1994). Clicnical cutoffs for the beck depression inventory and the geriatric depression scale witholder adult psychiatric outpatints. *Journal of Psychopathology and behavioral assessment, 16*, 3 233-242.

Kraepelin, E. (1913) Das manisch-depressive Irresein. In: Psychiatrie. Ein Lehrbuch fur Studierende und Urzte. Barth, Leipzig, pp 1183-1395

Kroenke K, Spitzer RL, Williams JBW. The patient health questionnaire-2: validity of a two-item depression screener. *Med Care.* 2003;41:1284-1292.

Kroenke K, Unützer J, Callahan CM, et al.

Patients With Major Depressive Disorder or Bipolar Disorder Am J Psychiatry 161:1433-1441, August 2004

Pirraglia, P. A., J. C. Peterson, et al. (1999). "Depressive symptomatology in coronary artery bypass graft surgery patients." Int J Geriatr Psychiatry 14(8): 668-80.

Wang, P. S., Beck, A. L., Berglund, P., McKenas, D. K., Pronk, N. P., Simon, G. E., &